arXiv:1511.00185v2 [math.DG] 6 Nov 2015

A SHORT NOTE ON ADDITIVE FUNCTIONS ON RIEMANNIAN CO-COMPACT COVERINGS

MINH KHA

Abstract. The main purpose of this note is to provide a topological approach to defining additive functions on Riemannian co-compact normal coverings.

1. Introduction

Let X be a connected smooth Riemannian manifold of dimension n equipped with an isometric, properly discontinuous, free, and co-compact action of a discrete deck group G. Notice that the deck group G is finitely generated due to the Švarc-Milnor lemma and hence, $\mathrm{Hom}(G, \mathbb{R})$ is finite dimensional. Furthermore, the orbit space $M := X/G$ is a compact Riemannian manifold when equipped with the metric pushed down from X. The action of an element $g \in G$ on $x \in X$ is denoted by $g \cdot x$. Let π be the covering map from X onto M. Thus, $\pi(g \cdot x) = \pi(x)$ for any $(g, x) \in G \times X$.

Following [6], we define the class of additive functions on the covering X as follows:

Definition 1.1.

- A real smooth function u on X is said to be **additive** if there is a homomorphism $\alpha : G \to \mathbb{R}$ such that
$$u(g \cdot x) = u(x) + \alpha(g), \quad \text{for all} \quad (g, x) \in G \times X.$$
We denote by $\mathcal{A}(X)$ the space of all additive functions on X.
- A map h from X to $\mathbb{R}^m$ ($m \in \mathbb{N}$) is called a vector-valued additive function on X if every component of h belongs to $\mathcal{A}(X)$.

We remark that additive functions on co-compact covers appeared in various results such as studying the structure of positive G-multiplicative type solutions [1, 6], describing the off-diagonal long time asymptotics of the heat kernel [3] and the Green's function asymptotics of periodic elliptic operators [2] on a noncompact abelian cover of a compact Riemannian manifold.

A direct construction of additive functions on X can be found in either [4, Section 3] or [6, Remark 2.6]. However, this construction depends on the choice of a fundamental domain for the base M in X. A more invariant approach to defining additive functions on covers was mentioned briefly in [1, 4]. Our goal in this note is to present the full details of this approach for any co-compact covering.

2. Additive functions on co-compact normal coverings

We begin with the following notion (see [1, 4]):

Definition 2.1. Let $H^1_{DR}(M), H^1_{DR}(X)$ be De Rham cohomologies of M and X, respectively. We denote by $\Omega^1(M;G)$ the image in $H^1_{DR}(M)$ of the set of all closed differential 1-forms ω on M (modulo the exact ones) such that their lifts ω to X are exact. In other words, $\Omega^1(M;G)$ is the kernel of the homomorphism

$$\pi^* : H^1_{DR}(M) \to H^1_{DR}(X),$$

where π^* is the induced homomorphism of the covering map $\pi : X \to M$.

By De Rham's theorem, $\Omega^1(M;G)$ is a finite dimensional vector space. Indeed, more is true:

Lemma 2.2. $\Omega^1(M,G) \cong \mathrm{Hom}(G,\mathbb{R})$.

Proof. By Hurewicz's theorem (see e.g., [5]), the homologies $H_1(M)$ and $H_1(X)$ are isomorphic to the abelianizations of the fundamental groups $\pi_1(M)$ and $\pi_1(X)$, respectively. Therefore, we can identify De Rham cohomologies $H^1_{DR}(M)$ and $H^1_{DR}(X)$ with $\mathrm{Hom}(\pi_1(M),\mathbb{R})$ and $\mathrm{Hom}(\pi_1(X),\mathbb{R})$, correspondingly. Since X is a normal covering of M, $\pi_1(X)$ is a normal subgroup of $\pi_1(M)$ and moreover, the sequence

$$0 \to \pi_1(X) \to \pi_1(M) \to G \to 0$$

is exact. Because $\mathrm{Hom}(\cdot,\mathbb{R})$ is a contravariant exact functor, we deduce the exactness of the following sequence of vector spaces:

$$0 \to \mathrm{Hom}(G,\mathbb{R}) \to H^1_{DR}(M) \to H^1_{DR}(X) \to 0.$$

Hence, $\Omega^1(M,G)$ is isomorphic to $\mathrm{Hom}(G,\mathbb{R})$. □

Fixing a base point $x_0 \in X$. For any closed 1-form ω on M such that its lift to X is exact, there exists a unique function $f_\omega \in C^\infty(X,\mathbb{R})$ such that $\pi^*\omega = df_\omega$ and $f_\omega(x_0) = 0$. Equivalently,

$$f_\omega(x) = \int_{x_0}^{x} \pi^*\omega, \quad \forall x \in X.$$

Lemma 2.3. *For such 1-form ω, we have:*

i) Fix any $g \in G$, then $f_\omega(g \cdot x) - f_\omega(x)$ is independent of $x \in X$.
ii) If $\pi^*\omega = 0$ then $\omega = 0$.

Proof.

i) For each $g \in G$, let L_g be the diffeomorphism of X that maps x to $g \cdot x$. Since $\pi \circ L_g = \pi$, we get $df_\omega = \pi^*\omega = L_g^*\pi^*\omega = L_g^* df_\omega = dL_g^* f_\omega = d(f_\omega \circ L_g)$. Thus, $d(f_\omega - f_\omega \circ L_g) = 0$ and so, $f_\omega \circ L_g - f_\omega$ is constant since X is connected.
ii) Fix any point $p \in M$. We pick an evenly covered open subset U of M such that it contains p. Then there is a smooth local section $\sigma : U \to X$, i.e., $\pi \circ \sigma = id_{|U}$ (see e.g., [5, Proposition 4.36]). Hence, $\omega(p) = \sigma^*\pi^*\omega(p) = 0$.

□

On $\mathcal{A}(X)$, we introduce an equivalent relation $\sim$ as follows: $f_1 \sim f_2$ in $\mathcal{A}(X)$ if and only if $f_1 - f_2 = f \circ \pi$ for some function $f \in C^\infty(M,\mathbb{R})$.

By Lemma 2.3 (i), the map $\omega \mapsto f_\omega$ induces the following linear map

$$\begin{aligned} \Lambda : \Omega^1(M,G) &\to \mathcal{A}(X)/\sim \\ [\omega] &\mapsto [f_\omega], \end{aligned} \tag{1}$$

where $[\omega]$, $[f_\omega]$ are the equivalent classes of ω, f_ω in $\Omega^1(M,G)$ and $\mathcal{A}(X)/\sim$, correspondingly. We now claim that $[\omega]=0$ if and only if $[f_\omega]=0$, and hence Λ is an injective linear map. Indeed, due to Lemma 2.3 (ii), the condition that ω is exact is equivalent to $\pi^*\omega = d(f\circ\pi)$ for some $f \in C^\infty(M,\mathbb{R})$. But this is the same as $df_\omega = d(f\circ\pi)$, or $[f_\omega]=0$.

Consider an additive function f on X. According to the definition, there exists a unique group homomorphism $\ell_f : G \to \mathbb{R}$ such that $f(g\cdot x) = f(x) + \ell_f(g)$ for any $g\in G$, $x\in X$. Then the map $f\mapsto \ell_f$ induces the linear map

$$\begin{aligned} \Upsilon : \mathcal{A}(X)/\sim &\to \mathrm{Hom}\,(G,\mathbb{R}) \\ [f] &\mapsto \ell_f, \end{aligned} \tag{2}$$

which is injective.

Then the composition $\Upsilon\circ\Lambda$ is also injective. By Lemma 2.2, $\dim_{\mathbb{R}}\Omega^1(M,G) = \dim_{\mathbb{R}}\mathrm{Hom}\,(G,\mathbb{R}) < \infty$. These facts together imply that the linear maps Υ and Λ are isomorphism. We conclude:

Theorem 2.4. *The three vector spaces $\Omega^1(M,G)$, $\mathcal{A}(X)/\sim$ and $\mathrm{Hom}\,(G,\mathbb{R})$ are isomorphic to each other.*

In particular, we obtain:

Corollary 2.5. *Assume that $G=\mathbb{Z}^d$. Then there is a smooth $\mathbb{R}^d$-valued function h on X such that for any $(g,x)\in\mathbb{Z}^d\times X$,*

$$h(g\cdot x) = h(x) + g. \tag{3}$$

The following standard proposition says that given any additive function u on X, then one can pick a harmonic additive function f such that $f-u$ is G-periodic.

Proposition 2.6. *For any ℓ in $\mathrm{Hom}(G,\mathbb{R})$, there exists a unique (modulo a real constant) **harmonic** function f on X such that for any $(g,x)\in G\times X$, we have*

$$f(g\cdot x) = f(x) + \ell(g). \tag{4}$$

Proof. First, we show the existence part. Due to Theorem 2.4, let $\tilde{f}$ be a function on X satisfying $\tilde{f}(g\cdot x) = \tilde{f}(x)+\ell(g)$ for any $(g,x)\in G\times X$. We recall the isomorphism Λ defined in (1). We put $\alpha := \Lambda^{-1}([\tilde{f}])\in\Omega^1(M,G)$. By the Hodge theorem, there exists a unique harmonic 1-form ω on M such that $[\omega]=\alpha$ in $H^1_{DR}(M)$. Let f be a smooth function such that $f\in\mathcal{A}(X)$ and $\pi^*\omega = df$. Then f satisfies (4) since $[f]=[\tilde{f}]$ in $\mathcal{A}(X)/\sim$. Thus, it is sufficient to show that f is harmonic on X. We denote by δ_X, Δ_X and δ_M, Δ_M the codifferential and Laplace-Beltrami operators on X and M, respectively. Since the covering map π is a local isometry between X and M, its pullback π^* intertwines the codifferential operators, i.e.,

$$\delta_X\pi^* = \pi^*\delta_M.$$

Since $\Delta_M \omega = 0$, it follows that $\delta_M \omega = 0$. Thus,

$$\Delta_X f = \delta_X df = \delta_X \pi^* \omega = \pi^* \delta_M \omega = 0.$$

For the uniqueness part, let f_1 and f_2 be any two harmonic functions on X such that (4) holds for each of these functions. Since $f_1 - f_2$ is G-periodic, it can be pushed down to a real function f on M. Moreover, $\pi^* \Delta_M f = \Delta_X \pi^* f = \Delta_X (f_1 - f_2) = 0$. Therefore, f must be constant since it is a harmonic function on a compact, connected Riemannian manifold M. Thus, $f_1 - f_2$ is constant. □

Corollary 2.7. *Fixing a base point x_0 in X. Then to each $\alpha \in \mathrm{Hom}(G, \mathbb{R})$, there exists a unique harmonic function f_α defined on X such that $f_\alpha(x_0) = 0$ and $\Upsilon([f_\alpha]) = \alpha$, where Υ is introduced in (2). Consequently,*

$$\mathcal{A}(X) = \bigsqcup_{\alpha \in \mathrm{Hom}(G, \mathbb{R})} \big\{ f_\alpha + \varphi \mid \varphi \text{ is periodic} \big\}.$$

Remark 2.8. When $G = \mathbb{Z}^d$, the Albanese pseudo-metric d_G introduced in [3, Section 2] is actually the pseudo-distance arising from **any** harmonic vector-valued additive function h satisfying (3), i.e., $d_G(x, y) = |h(x) - h(y)|$ for any $x, y \in X$.

3. Acknowledgements

The author was partially supported by the NSF grant DMS-1517938.

References

[1] S. Agmon, *On positive solutions of elliptic equations with periodic coefficients in $\mathbf{R}^n$, spectral results and extensions to elliptic operators on Riemannian manifolds*, Differential equations (Birmingham, Ala., 1983), 1984, pp. 7–17.

[2] M. Kha, *Green's function asymptotics of periodic elliptic operators on abelian coverings of compact manifolds*. arXiv:1511.00276, preprint.

[3] M. Kotani and T. Sunada, *Albanese maps and off diagonal long time asymptotics for the heat kernel*, Comm. Math. Phys. **209** (2000), no. 3, 633–670.

[4] P. Kuchment and Y. Pinchover, *Liouville theorems and spectral edge behavior on abelian coverings of compact manifolds*, Trans. Amer. Math. Soc. **359** (2007), no. 12, 5777–5815.

[5] J. M. Lee, *Introduction to smooth manifolds*, Second Edition, Graduate Texts in Mathematics, vol. 218, Springer, New York, 2013.

[6] V. Ya. Lin and Y. Pinchover, *Manifolds with group actions and elliptic operators*, Mem. Amer. Math. Soc. **112** (1994), no. 540, vi+78.

M.K., Department of Mathematics, Texas A&M University, College Station, TX 77843-3368, USA

E-mail address: kha@math.tamu.edu